

Description

FUNCTIONAL FOODS AND COSMETICS CONTAINING
SPHINGOGLYCOLIPIDS AND METHOD OF PRODUCING THE SAME

5

Technical Field:

This invention relates to functional foods and cosmetics containing sphingoglycolipids and a method of producing the same.

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000

Background Art:

Recent researches have been revealing that some complex lipids, particularly some glycolipids exhibit pronounced physiological activities. For example, it has come to be known that ceramides composed of sphingosine and fatty acids and cerebrosides composed of sphingosine, sugar, and fatty acids are present in the stratum corneum of human skin and serve to prevent evaporation loss of water from the body. Application of the glycolipids to the cosmetics taking advantage of this high moisturizing activity and application to pharmaceuticals taking advantage of elastase inhibitory activity or free radical inhibitory activity have advanced.

20

Ceramide related substances typified by the sphingoglycolipids have been supplied as extract from bovine brain, etc. However, since the BSE (bovine

25

spongiform encephalopathy) outbreak in 1986 the supply has decreased markedly because of the risk of human infection. The demand for safe ceramide related substances of plant origin has thus been increasing.

5 Seeing that ceramide related substances of plant origin are by no means inferior to those of animal origin and involve neither side effects nor toxicity as has recently been revealed, extraction of ceramide related substances from plant raw materials are now under intensive study. Sphingoglycolipids of plant origin, particularly glycosylceramides, that have hitherto been reported include those obtained from cereals and beans, such as rice (see *Agric. Biol. Chem.*, vol. 49, p. 2753 (1985)), rice bran (see JP-A-62-187404 and JP-A-11-279586), wheat (see *Agric. Biol. Chem.*, vol. 49, 1. 3609 (1985) and a published Japanese translation of a PCT patent application No. Hei.-6-507653), and soybeans (see *Chem. Pharm. Bull.*, vol. 38, No. 11, p. 2933 (1990)).

 Ceramides have been added to skin care products, hair care products, bathing preparations, and the like so that the manner of taking in ceramides has been almost restricted to percutaneous absorption. In recent years, it has been reported that one-month administration of 20 mg/day of wheat extract containing 3 wt% of wheat ceramides results in improvement on skin's moisture retention (see *Fragrance Journal*, vol. 23, No. 1, p. 81

(1995)), and oral intake of ceramides in the form of eatable cosmetics has been attracting attention.

Cosmetics containing ceramides of animal (horse) origin, ceramides produced by yeast fermentation or synthetic ceramides are now commercially available. Being safe, ceramides of plant origin have a favorable image, and many cosmetics featuring the plant origin are on the market.

However, plants which have been so far utilized as materials for supplying plant sphingoglycolipids are limited to cereals and beans, of which the sphingoglycolipids content is not much, usually about 0.01% by weight. All of these plant materials are edible for human consumption, whereas residue after extracting sphingoglycolipids therefrom has no value as food. The problem of using plants lies in that a vast quantity of food materials lose their value as food only to provide a slight amount of sphingoglycolipids. Looking out over the food processing industry, on the other hand, it is noted that 3,000 to 4,000 tons of "tobiko" powder occurs a year as a by-product in making konjac products (devil's tongue) from konjac (elephant foot or *Amorphophallus konjac*). Tobiko powder finds no use as a food due to its peculiar acrid taste and irritating smell, except that part of tobiko powder is made use of as a fertilizer or a concrete thickener. Further, cotton seed oil cake, which is a by-

product in squeezing cotton seed to obtain cotton seed oil, had been used as an animal feed rich in vegetable protein until 10 years ago. However, giving cow cotton seed as a high-energy feed is increasing among dairy farmers aiming to improve fat content in milk. Increase of utility of cotton seed oil cake is therefore demanded.

The present invention relates to functional foods and cosmetics containing sphingoglycolipids of plant origin which are attracting attention as a functional material of cosmetics and foods. It is an object of the present invention to provide functional foods and cosmetics made of materials of plant origin which are rid of the safety problem often pointed out in relation to the conventional materials extracted from animal tissue and yet find no value as food for the time being.

Disclosure of the Invention:

The present inventors have sought for plant raw materials containing sphingoglycolipids in a higher concentration than the conventionally used plant raw materials such as cereals and beans. As a result, they have unexpectedly found out that natural resources of plant origin, such as tuberous and corm vegetables (e.g., tobiko powder (fly powder) by-produced in making konjac (*Amorphophallus konjac*) products), and oil cakes (e.g., cotton seed oil cake), contain sphingoglycolipids in

concentrations comparable to or higher than those in cereal and bean crops and completed the present invention. More specifically, they have found that tuberous and corm vegetables including sweet potato and white potato as well as konjac tobiko powder, which occurs in quantity in making konjac products and is of little utility value as food, and oil cakes such as cotton seed oil cake are rich in sphingoglycolipids, particularly cerebrosides; that these components can be extracted efficiently from the above-mentioned natural resources by using organic solvents; and that foods or cosmetics containing the extract exhibit higher moisture retaining action than those containing conventional sphingoglycolipids of plant (cereal) origin. The present invention has been reached based on these findings.

A first aspect of the invention consists in functional foods characterized by containing sphingoglycolipids derived from tuberous and corm vegetables, preferably konjac, or oil cakes.

A second aspect of the invention consists in functional foods containing sphingoglycolipids and plant sterols. Preferably the sphingoglycolipids are those extracted from tuberous and corm vegetables, particularly konjac, or oil cakes.

A third aspect of the invention resides in cosmetics characterized by containing sphingoglycolipids

derived from tuberous and corm vegetables, preferably konjac, or oil cakes.

A fourth aspect of the invention lies in a method of producing sphingoglycolipids-containing products, functional foods or cosmetics characterized by comprising
5 adding an organic solvent to a tuberous and corm vegetable or oil cake and extracting sphingoglycolipids.

Best Mode for Carrying out the Invention:

10 The present invention will be hereinafter described in detail.

The plants which can be used as raw materials in the present invention include any kind of tuberous and corm vegetables and oil cakes, such as sweet potato, white
15 potato, taro, yam, konjac, long yam, cotton seed oil cake, rapeseed oil cake, coconut oil cake, and palm oil cake. They can be used either as such or after processed by drying, grinding, heating or a like operation. Preferred of them is konjac. Konjac tobiko powder is particularly
20 preferred because it is to be discarded in quantity and is easily available.

The sphingoglycolipids which can be used in the invention can be obtained by, for example, extraction from the above-recited plant raw materials with an organic
25 solvent as follows.

Any organic solvents can be used as an extracting solvent as long as they are not reactive on the raw material and sphingoglycolipids during extraction and do not impair the effects of the present invention. The solvents can be used either individually or as a mixture of two or more thereof. Such organic solvents include alcohols, such as methanol, ethanol, 1-propanol, 2-propanol, 1-butanol, 2-butanol, and tert-butanol; polyhydric alcohols, such as ethylene glycol, propylene glycol, butylene glycol, and glycerol; ketones, such as acetone and methyl ethyl ketone; esters, such as methyl acetate and ethyl acetate; ethers, such as tetrahydrofuran and diethyl ether; halogenated hydrocarbons, such as dichloromethane, dichloroethane, and chloroform; aliphatic hydrocarbons, such as hexane, pentane, and petroleum ether; aromatic hydrocarbons, such as toluene; polyethers, such as polyethylene glycol; and pyridines. Preferred of them are methanol, ethanol, acetone, and hexane. Methanol and ethanol are particularly preferred. In using these polar organic solvents for extraction, extraction efficiency may be improved by addition of water, additives such as surface active agents, and the like in amounts that do not impair the effects of the invention.

The amount of the organic solvent used for extraction is desirably about 1 to 30 times, particularly about 1 to 10 times, the plant raw material. With the

amount of the organic solvent being not smaller than this range, distribution of the solvent throughout the raw material is ensured to achieve sufficient extraction. Use of a greater amount of the solvent than this range results in no more improvement in extractability. The amount of the solvent falling within this range minimizes the load of solvent removal in the subsequent concentration step.

The extracting temperature depends on the boiling point of the solvent. In the case of methanol or ethanol, a suitable temperature is from room temperature up to 70°C, preferably from about room temperature to 60°C. At temperatures not lower than room temperature, the extraction efficiency can be increased. Raising the temperature over the above range is not so influential on extraction efficiency. Energy consumption can be reduced at temperatures not higher than the above range.

The extracting time is 10 minutes to 24 hours, preferably 1 to 10 hours. The extracting time being not shorter than 10 minutes, extraction can be carried out more sufficiently. Since an increase of extractability is not expected with a longer time than 24 hours, the above time range ensures completion of extraction in a minimized time.

The extraction operation is not limited to a single batch operation. A fresh solvent may be added to the raffinate to repeat extraction, or an extracting solvent may be brought into contact with the feed a plurality of

times. That is, the extraction operation can be carried out in any of a batch process, a semi-continuous process, and a countercurrent multistage process. Known extraction processes such as Soxhlet extraction may be used.

5 The extraction residue is then separated. Methods of separation are not particularly limited. Known methods, such as filtration by suction, pressure filtration using a filter press or a cylinder press, decantation, centrifugation, and centrifugal filtration, can be
10 employed.

 The extract thus obtained is preferably sent to a concentration step. The method of concentration is not particularly limited. For example, the extract is concentrated by means of a vacuum concentration apparatus,
15 such as an evaporator, or a centrifugal thin film evaporator, such as Evapol (from Ohkawara Seisakusho) or by heating to remove the solvent.

 The resulting concentrate can be used as such or, if necessary, be purified in a conventional manner to
20 remove impurities and increase the purity. Useful purification procedures include washing with water, washing with hexane, passing through a silica gel column, a resin column, a reversed-phase column, etc., partitioning between solvents having different polarities, and
25 recrystallization. Where obtaining highly pure sphingoglycolipids is particularly needed, purification is

preferably conducted by treating the concentrate with an alkaline solution, partitioning with chloroform, diethyl ether, etc., collecting and concentrating the organic layer, and separating the sphingoglycolipids by column chromatography on silica gel.

The sphingoglycolipids in the resulting concentrate are analyzed most conveniently by thin layer chromatography using commercially available sphingoglycolipids, especially glucosylceramide as a standard. The concentrate is applied on a silica gel thin layer plate and developed with an appropriate solvent system, such as a chloroform-methanol system. Visualization of the spots by color development with concentrated sulfuric acid or an anthrone reagent makes it easy to judge the presence of sphingoglycolipids in the concentrate in a high concentration. Presence of sphingoglycolipids in abundance can also be ascertained by other conventional techniques, such as high-performance liquid chromatography and various chromatography/mass spectrometries.

The functional foods as referred to herein are preparations which can be orally taken to exert one or more than one effects selected from skin moisture retention, improvement on skin roughening, skin beautification, and treatment of atopic dermatitis, allergic dermatitis, psoriasis, pimples, skin aging, hair loss, cancers, AIDS, hypertension, cholesterolemia, arteriosclerosis, and so on.

The concentrate may be taken directly as a functional food of the present invention but, for easy handling, is preferably mixed with carriers and/or diluents acceptable as food and drink additives and formed into preparation
5 forms, such as powders, tablets, capsules, gels, aqueous dispersions, ethanolic solutions or edible oil solutions.

Carriers which can be preferably used in powders, tablets or capsules include dextrin, cyclodextrins, potato starch, corn starch, and lactose. Carriers useful in gels
10 include gelatin, agar, guar gum, gum arabic, and konjac mannan. Water for use in aqueous dispersions is not limited. Water from springs or deep sea water may be used.

The aqueous dispersions can contain saccharides, amino acids, and inorganic salts as far as the present invention
15 is not ruined. Fruit juice or carbonated water is also useful. An emulsifier is preferably added to the aqueous dispersion to help the concentrate be effectively dispersed. Useful emulsifiers include sucrose fatty acid esters, glycerol fatty acid esters, polyglycerol fatty acid
20 esters, propylene glycol fatty acid esters, and soybean lecithin. Ethanol to be used in the ethanolic solutions is preferably distilled alcohol. The concentration of ethanolic solutions is not particularly limited. The edible oils to be used in the edible oil solutions are not
25 particularly limited and include soybean oil, rapeseed oil, olive oil, sesame oil, safflower oil, and wheat germ oil.

In order to effectively disperse the concentrate of the invention in edible oil, an emulsifier can be added. Suitable emulsifiers include soybean lecithin, sucrose fatty acid esters, glycerol fatty acid esters, polyglycerol fatty acid esters, and propylene glycol fatty acid esters.

The functional foods having the above-described preparation forms can be orally taken alone or as mixed into foods and/or beverages. The foods and/or beverages are not particularly limited and include, for example, staple foods such as bread, wheat noodle, buckwheat noodle, and rice; confectionery, such as cookies, cake, jelly, pudding, candy, chewing gum, and yogurt; liquors; and beverages, such as soft drinks, nutritious drinks, coffee, tea, and milk. In order to promote the effects, the mixed foods can further contain vitamins, collagen, squalane, soybean lecithin, plant sterols, hyaluronic acid, sorbitol, chitin, chitosan, glucosamine, niacin, niacinamide, Centella asiatica extract, polyphenols, co-enzyme Q10, pycnojenol, deep sea water, and the like. In particular, addition of plant sterols is preferred for producing high effects on skin moisture retention, atopic dermatitis, etc.

Any kind of plant sterols extracted from plants can be used in the invention. Specifically, waste such as deodorized scum oil occurring in production of oil, e.g., rice bran oil, cotton seed oil or soybean oil can be used. They are also extractable from konjac tobiko powder or the

waste from cane sugar factories. β -Sitosterol, campesterol, stigmasterol, sitostanol, campestanol, stigmastanol, and derivatives thereof can be used either individually or as a mixture of two or more thereof. Any derivatives, either synthesized or naturally-occurring, can be used unless they impair the effects of the invention. Sterol glycosides having sugar bonded, esterified compounds, and compounds having an amide linkage are preferred.

The plant sterols used in the invention can be obtained by known processes. A typical process comprises alkali treatment of deodorized scum oil which is by-produced in the deodorization step of soybean oil by steam distillation, extraction with an organic solvent, e.g., hexane, and crystallization from a poor solvent, e.g., acetone.

Commercially available plant sterols for foods as supplied by Eisai Co., Ltd. (trade name: Phytosterol F), Tuno Food Industrial Co., Ltd., etc. are also employable in the present invention.

Where plant sterols are taken in combination with sphingoglycolipids according to the present invention, the sphingoglycolipids to be combined may be any of synthetic products, fermentation products, and products of animal or plant origin. From the standpoint of use as a food, those of plant origin which are of high safety and have been

eaten are preferred. Any sphingoglycolipids of plant origin can be used as far as they are extracted from edible plants. Preferred are those extracted from cereal crops, such as wheat, wheat bran, barley, rice, rice bran, and corn; beans, such as soybeans; greens, such as spinach, komatuna, and boy choy; tuberous and corm vegetables, such as sweet potato, white potato, taro, yam, konjac, and long yam; and oil cakes, such as cotton seed oil cake, rapeseed oil cake, coconut oil cake, and palm oil cake. Those extracted from the tuberous and corm vegetables or oil cakes are still preferred. In particular, sphingoglycolipids extracted from konjac are the most preferred for their high effects.

Where sphingoglycolipids and plant sterols are mixed into other foods and/or beverages, their contents in the mixtures vary according to the food or food intake and cannot be generally specified but may be represented in terms of their intakes. An advisable daily intake of sphingoglycolipids is 1 μ g to 100 g, and an advisable daily intake of plant sterols is 1 mg to 1000 g. The contents in foods are adjusted so as to give such levels of intake.

The material containing sphingoglycolipids extracted from plants can be powdered by drying by lyophilization, spray drying, vacuum drying or a like technique. If necessary, the dried product is ground in a sample mill, a speed mill, a blender, a mixer, etc. The

resulting powder may sometimes exhibit viscosity. In such cases, carriers capable of accelerating powderization can be added to such an extent that would not damage the effects of the invention. Such carriers include corn
5 starch, potato starch, dextrin, cyclodextrins, wheat flour, bread crumbs, edible salt, zeolite, talc, and oyster shell powder.

Tablets are prepared by putting the powder preparation obtained as described above in a tablet machine and punching into tablets. A binder may be used in such an amount that does not impair the effects of the invention.

Capsules are prepared by putting the powder preparation obtained as described above and/or a solution preparation into capsules for medicines or foods. Any
15 capsules can be used unless the effects of the invention are impaired.

Gels are prepared by any known technique as long as the effects of the present invention are not ruined. For example, the resulting sphingoglycolipids-containing
20 product is dispersed and/or dissolved in water, and a conventionally known gelling agent, thickening polysaccharide, etc. is added thereto. Gelling agents used for foods, such as gelatin and agar, are used preferably.

Aqueous dispersions are prepared by stirring the
25 sphingoglycolipids-containing product in a prescribed amount of water. Dispersing may be accelerated by

subjecting the system to ultrasonication or adding a known emulsifier or dispersant as long as the effects of the present invention are not damaged.

Soluble in ethanol, the sphingoglycolipids-
5 containing product is simply dissolved in ethanol to give an ethanolic solution. If desired, a stabilizer may be added to the ethanolic solution to prevent precipitation or turbidity within such an amount that does not impair the effects of the invention.

10 Soluble in edible oils, the sphingoglycolipids-
containing product can be directly dissolved in edible oils to provide edible oil solutions. Where difficult to dissolve, the sphingoglycolipids-containing product is once
15 dissolved in a small amount of soybean lecithin or egg yolk lecithin and then dissolved in an edible oil to prepare an edible oil solution with ease. In order to prevent precipitation or turbidity, a stabilizer, etc. may be added
20 within such an amount that does not impair the effects of the invention. Further, dissolving can be accelerated by slightly heating the system to reduce the viscosity of the edible oil.

The edible oils which can be used in the functional foods of the invention are not particularly limited. Useful edible oils include animal fats and oils, such as
25 lard, beef tallow, whale oil, fish oil, and butter; vegetable oils, such as rapeseed oil, soybean oil, olive

oil, rice oil, sunflower oil, and coconut oil; and functionalized synthetic oils or mixed oils. They can be used either individually or as a mixture of two or more thereof. Preferred of them are vegetable oils and health-
5 friendly synthetic oils.

Made up of components of plant origin, the functional foods according to the present invention have high safety. Therefore, the oral intake and the interval of intake are not particularly limited. An advisable daily
10 intake for manifestation of the aimed effects of the invention is from 1 μ g to 100 g in terms of the sphingoglycolipids content. It is recommended to maintain taking at a frequency of ten times a day to once per three days. It is advisable to continue taking for at least 7
15 days, preferably 14 days or longer.

The cosmetics as referred to in the present invention include preparations which, when applied to the skin or hair, penetrate inside to exert one or more than one effects selected from skin moisture retention, skin
20 whitening, improvements on wrinkles, dark blotches, freckles, and skin roughening, hair growth, prevention of hair splitting, and treatment of atopic dermatitis, allergic dermatitis, pimples, xeroderma, and the like. Accordingly, the present invention embraces in its scope a
25 method for treating and/or preventing these symptoms which comprises applying the sphingoglycolipids of the invention

in an amount effective on treatment and/or prevention of these symptoms. The formula may solely comprise the sphingoglycolipids or be a combination with other cosmetic materials, i.e., cosmetically acceptable carriers and/or diluents.

Cosmetically acceptable carriers and/or diluents are not particularly limited. Examples include colorants, such as titanium mica, talc, silica, and titanium dioxide; oily materials, such as olive oil, castor oil, jojoba oil, bees wax, lanolin, squalane, cetanol, liquid paraffin, methylpolysiloxane, silicone polyether copolymers, and behenyl alcohol; humectants, such as glycerol, hyaluronic acid, sorbitol, propylene glycol, dipropylene glycol, chitin, and chitosan; surface active agents, such as alkylsulfates and alkyl ether sulfates; emulsifiers, such as sucrose fatty acid esters, polyglycerol fatty acid esters, sorbitan fatty acid esters, lecithin, polyoxyethylene alkyl ethers, and alkyl glucosides; vegetable gum or water-soluble polymers, such as quince seed gum, xanthan gum, sodium carboxymethyl cellulose (CMC), carboxyvinyl polymers, polyvinylpyrrolidone, and polyvinyl alcohol; water; and ethanol.

The form is not restricted. For example, the cosmetics can be applied to the skin in the form of lotions, emulsions, moisture creams, sunscreens, anti-sunburn cosmetics, face packs, foundations, finishing

powders, rouge, eye make-ups, perfumes, eau-de-colognes, lip creams, lipsticks, etc.; or to the hair in the form of hair growth products, pomade, setting lotions, hair sprays, hair dyes, hair tonics, eyelash cosmetics, etc. The
5 formula can be added to facial cleansing creams, facial soaps, shampoos, hair conditioners, hair treatments, bathing preparations, and the like.

The method for extracting sphingoglycolipids to be incorporated into the cosmetics of the invention is the
10 same as for extracting sphingoglycolipids to be incorporated into the functional foods. The sphingoglycolipids as extracted may be applied directly as cosmetics of the invention. For easier handling, the extract is dissolved in an organic solvent such as ethanol
15 or dispersed in water.

Soluble in ethanol, the sphingoglycolipids-containing product is simply dissolved in ethanol to give an ethanolic solution. A stabilizer may be added to the solution to prevent precipitation or turbidity within such
20 an amount that does not impair the effects of the invention.

The aqueous dispersion is prepared by stirring the sphingoglycolipids-containing product in a prescribed amount of water. Dispersing may be accelerated by
25 subjecting the system to heat treatment or ultrasonication or adding a known emulsifier or dispersant as long as the

effects of the present invention are not damaged. In particular, it is preferred that the sphingoglycolipids-containing product be once dissolved in an emulsifier, such as soybean or egg yolk lecithin, hydrogenated lecithin and then dispersed in water.

In order to enhance the effects, the cosmetics of the invention can further contain vitamins, collagen, squalane, soybean lecithin, plant sterols, hyaluronic acid, sorbitol, chitin, chitosan, glycerol, butylene glycol, propylene glycol, niacin, niacinamide, galactosylceramide, and so forth.

Containing the sphingoglycolipids derived from tuberous and corm vegetables or oil cakes, the cosmetics according to the present invention are of high safety and may have an unlimited sphingoglycolipids content. An advisable content of the sphingoglycolipids from tuberous and corm vegetables or oil cakes for manifestation of the aimed effects of the invention is from 0.001% by weight to 100% by weight. At a content not smaller than 0.001 wt%, the effects of the invention show up sooner and more sufficiently.

The typical effects produced when the functional foods or cosmetics of the invention are taken orally or applied to the skin are enhancement of skin moisture retention and improvement on skin roughening. With respect to the skin moisture retention, the effect can be

ascertained through known means such as measurement of transepidermal water loss (TEWL) or measurement of water content of the stratum corneum (hereinafter referred to as an SC water content). TEWL measurement can be made with, for example, Evaporimeter from Servo Med, Sweden or Tewameter from Courage + Khazaka Electronic GmbH, German. Measurement of the SC water content can be taken with, for example, Corneometer from Courage + Khazaka, German or Skikon 200 from IBS. The effect on improvement of skin roughening can be seen by observation with the naked eye and by the user's own feel. The skin whitening effect can also be seen by observation with the naked eye or quantitatively measured with a skin color meter Mexameter MX16 from Courage + Khazaka, German.

The effects produced by the functional foods or cosmetics of the invention orally taken or applied to the skin owe to the specific actions on the skin of the sphingoglycolipids contained in tuberous and corn vegetables or oil cakes. Therefore, the functional foods and cosmetics of the invention are superior to those containing the sphingoglycolipids obtained from conventional plant materials, such as wheat, rice bran, and soybeans, particularly in improving effects on skin roughening, atopic dermatitis, allergic dermatitis, and pimples.

Examples:

The present invention will now be illustrated in greater detail with reference to Examples, but the present invention is not deemed to be limited thereto. To begin with, the measuring apparatus and methods used in Examples are described.

(1) Qualification of sphingoglycolipids

Sphingoglycolipids were qualitatively analyzed by silica gel thin layer chromatography (TLC). A prescribed amount of a sample was applied to a silica gel plate (Silicagel 60F254 Type from Merc; layer thickness: 0.5 mm), and the plate was placed in a developing chamber containing chloroform:methanol:water (=87:13:2 by volume) to develop the sample. After the development, the silica gel plate was dried with a drier, etc., and sulfuric acid was sprayed thereto followed by heating to cause color development.

(2) Quantification of sphingoglycolipids

Sphingoglycolipids were quantitatively analyzed by high-performance liquid chromatography (HPLC). HPLC was carried out by using LC Module 1 (from Waters Corp.), a column Inertsil SIL 100A (from GL Science Inc.), and chloroform:methanol (=9:1 by volume) as a solvent. The measurement was carried out at a flow rate of 1.0 ml/min at 37°C. A scattered light detector 500 ELSD from Alltech Associates Inc. was used for detection.

(3) Measurement of stratus corneum (SC) water content

Corneometer CM825 from Courage + Khazaka was used for SC water content measurement. Measurement was always made on the same site 1 cm below the left eye. Measurement
5 was taken 10 times for each test person to obtain an average value, which was taken as the datum of that person, and the SC water content was represented as an average of the data from all test persons.

10 Example 1: Extraction of sphingoglycolipids from konjac tobiko powder

One kilogram of konjac tobiko powder was put into a stirring tank, and 2 liters of ethanol was added, followed by stirring at room temperature for 2 hours. The mixture
15 was filtered to separate into an extract and a residue. The extract was concentrated in an evaporator to give 10.7 g of a brown waxy concentrate. The product was analyzed by the above-described methods for qualification and quantification. TLC revealed spots corresponding to
20 sphingoglycolipids. The amount of the sphingoglycolipids in the extract was found to be 0.55 g by HPLC. The purity of the sphingoglycolipids in the tobiko powder extract was found to be 5.1 wt%.

Example 2: Extraction of sphingoglycolipids from cotton seed oil cake

One kilogram of cotton seed oil cake was put into a stirring tank, and 2 liters of ethanol was added, followed by stirring at room temperature for 5 hours. The mixture was filtered to separate into an extract and a residue. The extract was concentrated in an evaporator to give 8.4 g of a brown waxy concentrate. The product was analyzed by the above-described methods for qualification and quantification. TLC revealed spots corresponding to sphingoglycolipids. The amount of the sphingoglycolipids in the extract was found to be 0.41 g by HPLC. The purity of the sphingoglycolipids in the cotton seed oil cake extract was found to be 4.9 wt%.

Example 3: Extraction of sphingoglycolipids from potato peel

One kilogram of potato peel grinds (100 μ m or smaller particles) was put into a stirring tank, and 2 liters of ethanol was added, followed by stirring at room temperature for 2 hours. The mixture was filtered to separate into an extract and a residue. The extract was concentrated in an evaporator to give 10.2 g of a yellow waxy concentrate. The product was analyzed by the above-described methods for qualification and quantification. TLC revealed spots corresponding to sphingoglycolipids.

The amount of the sphingoglycolipids in the extract was found to be 0.42 g by HPLC. The purity of the sphingoglycolipids in the potato peel extract was 4.1 wt%.

5 Comparative Example 1: Extraction of sphingoglycolipids from wheat flour

One kilogram of wheat flour was put into a stirring tank, and 2 liters of ethanol was added, followed by stirring at room temperature for 5 hours. The mixture was filtered to separate into an extract and a residue. The extract was concentrated in an evaporator to give 6.8 g of a deep yellowish brown waxy concentrate. The product was analyzed by the above-described methods for qualification and quantification. The spots of sphingoglycolipids appearing in TLC were light in color. The amount of the sphingoglycolipids in the extract was found to be 0.05 g by HPLC. The purity of the sphingoglycolipids in the wheat flour extract was as low as 0.7 wt%.

20 Comparative Example 2: Extraction of sphingoglycolipids from rice bran

One kilogram of defatted rice bran was put into a stirring tank, and 3 liters of ethanol was added, followed by stirring at room temperature for 2 hours. The mixture was filtered to separate into an extract and a residue. The extract was concentrated in an evaporator to give

22.3 g of a brown waxy concentrate. The product was analyzed by the above-described methods for qualification and quantification. As a result of TLC, the spots of sphingoglycolipids appearing in TLC were light in color, while the spots corresponding to glyceroglycolipids, sterols, etc. developed deep colors. The amount of the sphingoglycolipids in the extract was found to be 0.38 g by HPLC. The purity of the sphingoglycolipids in the defatted rice bran extract was 1.7 wt%.

As described above, the content of sphingoglycolipid components in the tobiko powder extract obtained by using an organic solvent is as high as 5.1 wt% based on the extract and 0.055 wt% based on the feed. It has now been proved that sphingoglycolipid components are present in a tobiko powder extract in extremely high concentration and purity. It is also seen that sphingoglycolipid components are present in a potato peel extract and a cotton seed oil cake extract in sufficiently high concentration and purity as compared with the conventionally employed feeds for extraction, i.e., wheat flour and rice bran. Thus, the present invention is successful in recovering sphingoglycolipids from tuberous and corm vegetables and oil cakes in extremely high concentrations.

Example 4: Preparation of powdered functional food

In 20.0 g of ethanol was dissolved 10.0 g of the konjac tobiko powder extract obtained in Example 1, and the solution was introduced into 200 g of water while stirring.

- 5 The mixture was further stirred while in a dispersed state. Thirty minutes later, 5.0 g of sodium chloride was added, and the stirring was continued for an additional 10 minute period to give a brown precipitate, which weighed 6.8 g. The sphingoglycolipids content was 0.45 g as measured by HPLC. The extract was washed with water to have the sphingoglycolipids purity increased to 6.6 wt%. The washing operation was repeated once more to give 6.4 g of a precipitate (designated precipitate A) containing sphingoglycolipids of konjac tobiko powder origin. The sphingoglycolipids content was found to be 0.44 g as a result of HPLC. The purity of the sphingoglycolipids increased up to 6.9 wt% by washing twice.

- In 15 ml of ethanol was dissolved 5.0 g of precipitate A, and the resulting ethanolic solution was poured into a suspension of 15.0 g of corn starch in 50 ml of water while stirring, and the suspension was heated in a boiling water bath for 5 minutes while stirring. As the corn starch was dissolved and swollen, the suspension increased its viscosity and became pasty. The resulting pasty composition was dried in a vacuum drier at 50°C for 12 hours to give a milky white solid, which was ground in a

blender to obtain 19.3 g of a powdered functional food containing sphingoglycolipids of konjac tobiko powder origin. The resulting powder was soaked in ethanol, and the soluble components thus dissolved out were analyzed by
5 TLC. Spots of sphingoglycolipids were clearly observed.

Example 5: Preparation of powdered functional food

In 15 ml of ethanol was dissolved 5.0 g of precipitate A of Example 4. The resulting ethanolic
10 solution was poured into a suspension of 10.0 g of β -cyclodextrin in 20 ml of water while stirring. As stirring was continued, the suspension increased its viscosity and became pasty. The resulting pasty composition was dried in a vacuum drier at 50°C for 12 hours to give a milky white
15 solid, which was ground in a blender to obtain 14.8 g of a powdered functional food containing sphingoglycolipids of konjac tobiko powder origin. The resulting powder was soaked in ethanol, and the soluble components thus dissolved out were analyzed by TLC. Spots of
20 sphingoglycolipids were clearly observed.

Example 6: Preparation of powdered functional food

In 15 ml of ethanol was dissolved 5.0 g of precipitate A of Example 4. The resulting ethanolic
25 solution was poured into a suspension of 15.0 g of corn starch in 50 ml of water while stirring, and the suspension

was heated in a boiling water bath for 5 minutes while stirring. As the corn starch was dissolved and swollen, the suspension increased its viscosity and became pasty. To the resulting pasty composition was added 450 ml of water, followed by stirring to obtain a low-viscosity dispersion. The dispersion was delivered to a spray drier (Spray Dryer LT-8, manufactured by Ohkawara Kakoki K.K.) at a rate of 3 l/min and powdered under conditions of an inlet temperature of 250°C, an outlet temperature of 150°C, and an atomizer revolution speed of 35000 rpm. The resulting powdered functional food containing sphingoglycolipids of konjac tobiko powder origin weighed 18.7 g and assumed a light brown color. The resulting powder was soaked in ethanol, and the soluble components thus dissolved out were analyzed by TLC. Spots of sphingoglycolipids were clearly observed.

Example 7: Preparation of functional food in aqueous dispersion form

In 500 ml of water were put 5.0 g of precipitate A of Example 4 and 2.5 g of decaglycerol monostearate (SY Glyster MSW-750, available from Sakamoto Yakuhin Kogyo Co., Ltd.) and dispersed by ultrasonication using BRANSON 3200, available from Yamato Scientific Co., Ltd., for 30 minutes. There was obtained a functional food of aqueous dispersion form containing sphingoglycolipids of konjac tobiko powder

origin. The resulting aqueous dispersion was uniform and generated no sediment on 3 months' standing.

Example 8: Preparation of gelatinous functional food

5 In 15 ml of ethanol was dissolved 5.0 g of precipitate A of Example 4. The resulting ethanolic solution was poured into a solution of 15.0 g of gelatin in 20 ml of water heated at 60°C while stirring. The solution was allowed to cool to room temperature whereupon it turned into an ocherous hard gel. There was thus obtained a gelatinous functional food containing sphingoglycolipids of konjac tobiko powder origin.

Comparative Example 3: Preparation of food containing sphingoglycolipids of wheat flour origin

10 In 10.0 g of ethanol was dissolved 5.0 g of the wheat flour extract obtained in Comparative Example 1, and the solution was introduced into 100 g of water while stirring. The mixture was further stirred while in a dispersed state. Thirty minutes later, 2.5 g of sodium chloride was added, and the stirring was continued for an additional 10 minute period to give a brown precipitate, which weighed 3.6 g. The sphingoglycolipids content was 0.04 g as measured by HPLC. After washing with water, the
15
20
25 extract had a sphingoglycolipids purity of 1.1 wt%. The washing operation was repeated once more to give 3.4 g of a

precipitate containing sphingoglycolipids of wheat flour origin. The sphingoglycolipids content was found to be 0.04 g as a result of HPLC. The purity of the sphingoglycolipids obtained after washing twice was 1.2 wt%.

In 300 ml of water were put 3.0 g of the precipitate and 1.5 g of decaglycerol monostearate (SY Glyster MSW-750, available from Sakamoto Yakuhin Kogyo Co., Ltd.), and the mixture was treated in the same manner as in Example 7 to obtain an aqueous dispersion containing sphingoglycolipids of wheat flour origin.

Comparative Example 4: Preparation of food containing sphingoglycolipids of rice bran origin

In 20.0 g of ethanol was dissolved 10.0 g of the rice bran extract obtained in Comparative Example 2, and the solution was introduced into 200 g of water while stirring. The mixture was further stirred while in a dispersed state. Thirty minutes later, 5.0 g of sodium chloride was added, and the stirring was continued for 10 minutes to give a brown precipitate, which weighed 7.4 g. The sphingoglycolipids content was 0.24 g as measured by HPLC. After washing with water, the extract had a sphingoglycolipids purity of 3.2 wt%. The washing operation was repeated once more to give 6.9 g of a precipitate containing sphingoglycolipids of rice bran

origin. The sphingoglycolipids content was found to be 0.22 g as a result of HPLC. The purity of the sphingoglycolipids obtained after washing twice was 3.2 wt%.

5 In 300 ml of water were put 3.0 g of the precipitate and 1.5 g of decaglycerol monostearate (SY Glyster MSW-750, available from Sakamoto Yakuhin Kogyo Co., Ltd.), and the mixture was treated in the same manner as in Example 7 to obtain an aqueous dispersion containing
10 sphingoglycolipids of rice bran origin.

Test Example 1: Influences on SC water content

The functional food of the present invention obtained in Example 7 (aqueous dispersion), the aqueous
15 dispersions obtained in Comparative Examples 3 and 4, and a dispersion containing no sphingoglycolipids were tested for SC water content improving effects on ten volunteers for each sex (five persons between the ages of 20 and 40 and five persons between the ages of 40 and 60 for each sex).

20 The dispersion containing no sphingoglycolipids was prepared by dissolving 1.5 g of decaglycerol monostearate (SY Glyster MSW-750, available from Sakamoto Yakuhin Kogyo Co., Ltd.) in 300 ml of water.

The test persons were asked to take in 1.45 ml/day
25 of the functional food of the invention, 8.5 ml/day of the dispersion obtained in Comparative Example 3, and

3.25 ml/day of the dispersion obtained in Comparative Example 4 each for 30 days. These daily intakes correspond to 1 mg of the sphingoglycolipids per day. The dispersion containing no sphingoglycolipids as a control was taken in an amount of 1.45 ml/day.

The SC water content of the skin 1 cm below the left eye was measured with Corneometer after 10 days, 20 days and 30 days from the first day of taking. The average results of the twenty test persons are shown in Table 1.

TABLE 1

	Initial Value	After 10 Days	After 20 Days	After 30 Days
Example 7 (tobiko powder origin)	44	62	68	73
Comparative Example 3 (wheat flour origin)	47	51	55	58
Comparative Example 4 (rice bran origin)	46	49	53	60
Control	43	45	47	46

It is seen from Table 1 that taking sphingoglycolipids of konjac tobiko powder origin produces greater effects in improving the skin SC water content than taking sphingoglycolipids of wheat flour or rice bran origin.

Test Example 5: Improving effect on skin roughening

The functional food of the present invention obtained in Example 7 (aqueous dispersion), the aqueous dispersions obtained in Comparative Examples 3 and 4, and a dispersion containing no sphingoglycolipids were tested for rough skin improving effects with cooperation of ten females suffering from skin roughening on the back of their hands. The dispersion containing no sphingoglycolipids was prepared by dissolving 1.5 g of decaglycerol monostearate (SY Glyster MSW-750, available from Sakamoto Yakuhin Kogyo Co., Ltd.) in 300 ml of water.

The test persons were asked to take in 1.45 ml/day of the functional food of the invention, 8.5 ml/day of the dispersion obtained in Comparative Example 3, and 3.25 ml/day of the dispersion obtained in Comparative Example 4 each for 15 days. These daily intakes correspond to 1 mg of the sphingoglycolipids per day. The dispersion containing no sphingoglycolipids as a control was taken in an amount of 1.45 ml/day.

Meanwhile they were asked to answer a questionnaire about the condition of the back of their hands after 5 days, 10 days and 15 days. The results obtained are shown in Table 2 as a total score.

TABLE 2

	After 5 Days	After 10 Days	After 15 Days
Example 7 (tobiko powder origin)	22	29	34
Comparative Example 3 (wheat flour origin)	19	22	29
Comparative Example 4 (rice bran origin)	18	22	27
Control	8	13	14

Answers and scores:

4 ... The skin roughening was completely healed.

3 ... The skin condition was improved greatly.

2 ... The skin condition was improved.

1 ... The skin condition was unchanged.

0 ... The skin condition got worse.

It is seen from Table 2 that taking

sphingoglycolipids of konjac tobiko powder origin brings about more excellent effects of improving skin roughening than taking sphingoglycolipids of wheat flour or rice bran origin.

Example 9: Preparation of powdered functional food containing plant sterols

Plant sterols of soybean oil origin which mainly comprise β -sitosterol (2.0 g; Phytosterol F from Eisai Co., Ltd., hereinafter simply referred to as Phytosterol F) was

added to 14.8 g of the powder containing sphingoglycolipids of konjac tobiko powder origin obtained in Example 5, and the mixture was stirred well to get uniform to prepare a powdered functional food.

5

Example 10: Preparation of plant sterols-containing functional food of aqueous dispersion form

Precipitate A of Example 4 (5.0 g) and 2.0 g of Phytosterol F were kneaded at 50°C for 5 minutes to obtain a brown paste. The paste was put into 500 ml of water together with 2.5 g of decaglycerol monostearate (SY Glyster MSW-750, available from Sakamoto Yakuhin Kogyo Co., Ltd.) and dispersed by ultrasonication using BRANSON 3200, available from Yamato Scientific Co., Ltd., for 30 minutes. The resulting aqueous dispersion containing sphingoglycolipids and plant sterols was uniform and formed no sediment on 3 months' standing.

Example 11: Preparation of plant sterols-containing functional food of aqueous dispersion form

Precipitate A of Example 4 (1.0 g) and 2.0 g of Phytosterol F were kneaded at 50°C for 5 minutes to obtain a brown paste. The paste was put into 500 ml of water together with 2.5 g of decaglycerol monostearate (SY Glyster MSW-750, available from Sakamoto Yakuhin Kogyo Co., Ltd.) and dispersed by ultrasonication using BRANSON 3200,

available from Yamato Scientific Co., Ltd., for 30 minutes.

The resulting aqueous dispersion containing sphingoglycolipids and plant sterols was uniform and generated no sediment on 3 months' standing.

5

Example 12: Preparation of plant sterols-containing functional food of aqueous dispersion form

An aqueous dispersion containing sphingoglycolipids and plant sterols was prepared in the same manner as in Example 10, except for replacing Phytosterol F with β -sitostanol (from Sigma). The resulting aqueous dispersion containing sphingoglycolipids and plant sterols was uniform and generated no sediment on 3 months' standing.

10

15

Example 13: Preparation of plant sterols-containing gelatinous functional food

In 15 ml of ethanol was dissolved 5.0 g of precipitate A of Example 4, and 2.0 g of Phytosterol F was added thereto, followed by stirring at 30°C for 5 minutes to obtain a solution. The resulting ethanolic solution was introduced into an aqueous solution of 15.0 g of gelatin in 20 ml of water heated at 60°C while stirring. The solution was allowed to cool to room temperature whereupon it turned into an ocherous hard gel to give a gelatinous functional food.

20

25

Example 14: Preparation of plant sterols-containing wheat flour extract

The wheat flour extract (3.0 g) obtained in Comparative Example 1 and 0.21 g of Phytosterol F were kneaded at 50°C for 5 minutes to obtain an ochreous paste. The paste was put into 300 ml of water together with 1.5 g of decaglycerol monostearate (SY Glyster MSW-750, available from Sakamoto Yakuhin Kogyo Co., Ltd.). The mixture was treated in the same manner as in Example 10 to obtain an aqueous dispersion containing sphingoglycolipids of wheat flour origin and plant sterols.

Example 15: Preparation of plant sterols-containing rice bran extract

The rice bran extract (3.0 g) obtained in Comparative Example 2 and 0.56 g of Phytosterol F were kneaded at 50°C for 5 minutes to obtain a brown paste. The paste was put into 300 ml of water together with 1.5 g of decaglycerol monostearate (SY Glyster MSW-750, available from Sakamoto Yakuhin Kogyo Co., Ltd.). The mixture was treated in the same manner as in Example 10 to obtain an aqueous dispersion containing sphingoglycolipids of rice bran origin and plant sterols.

Comparative Examples 5 to 7

For comparison, an aqueous dispersion containing sphingoglycolipids of konjac tobiko powder origin was prepared in the same manner as in Example 10, except that β -sitosterol was not added (Comparative Example 5). In 500 g of water was dispersed 2.0 g of Phytosterol F with the aid of 2.5 g of decaglycerol monostearate (SY Glyster MSW-750, available from Sakamoto Yakuhin Kogyo Co., Ltd.) as an emulsifier to prepare an aqueous dispersion (Comparative Example 6). An aqueous solution of 2.5 g of the same decaglycerol monostearate in 500 g of water was prepared (Comparative Example 7).

Test Example 3: Influences on SC water content

Ten volunteers for each sex (five persons between the ages of 20 and 40 and five persons between the ages of 40 and 60 for each sex) were asked to take a prescribed amount of each functional food obtained in Examples for 30 days each. The SC water content of the skin 1 cm below the left eye was measured with Corneometer after 10 days, 20 days and 30 days. The functional foods and intakes used in the test are shown in Table 3.

The average results of the twenty test persons are shown in Table 3. Table 3 proves that taking the functional foods comprising the sphingoglycolipids of

konjac origin and plant sterols produces excellent effects
in improving the skin SC water content.

10088301-031802

TABLE 3

	Daily Intake			SC Water Content			
	Food	Sphingo- glycolipids	Plant Sterols	Initial Value	After 10 Days	After 20 Days	After 30 Days
Example 10	aqueous dispersion 1.45 ml	konjac 1 mg	Phytosterol F 5.8 mg	40	57	69	72
Example 11	aqueous dispersion 1.45 ml	konjac 0.2 mg	Phytosterol F 5.8 mg	43	58	66	69
Example 12	aqueous dispersion 1.45 ml	konjac 1.0 mg	β -sitostanol 5.8 mg	39	63	70	73
Example 14	aqueous dispersion 8.33 ml	wheat 1.0 mg	Phytosterol F 5.8 mg	42	56	61	64
Example 15	aqueous dispersion 3.13 ml	rice bran 1.0 mg	Phytosterol F 5.8 mg	40	55	63	66
Compara. Example 5	aqueous dispersion 1.45 ml	konjac 1.0 mg	-	41	55	60	63
Compara. Example 6	aqueous dispersion 1.45 ml	-	Phytosterol F 5.8 mg	39	42	43	45
Compara. Example 7	solution 1.45 ml	-	-	38	39	43	40

Phytosterol F: Plant sterols of soybean oil origin mainly comprising β -sitosterol, available from Eisai Co., Ltd.

Test Example 4: Skin roughening improving effects

Cooperation of ten females suffering from skin roughening on the back of their hands was obtained. The test persons were asked to daily take in a prescribed amount of each functional food prepared in Examples for 15 days. Meanwhile they were asked to answer a questionnaire about the condition of the back of their hands after 5 days, 10 days and 15 days. The samples and intakes used are shown in Table 4.

The results obtained are shown in Table 4 as a total score. It is seen from Table 4 that taking functional foods comprising the sphingoglycolipids of konjac origin and plant sterols produces excellent improving effects on skin roughening.

TABLE 4

	Daily Intake		Skin Roughening Improving Effects			
	Food	Sphingo-glycolipids	Plant Sterols	After 5 Days	After 10 Days	After 15 Days
Example 10	aqueous dispersion 1.45 ml	konjac 1 mg	Phytosterol F 5.8 mg	27	30	35
Example 11	aqueous dispersion 1.45 ml	konjac 0.2 mg	Phytosterol F 5.8 mg	22	28	32
Example 12	aqueous dispersion 1.45 ml	konjac 1.0 mg	β -sitosterol 5.8 mg	25	30	36
Example 14	aqueous dispersion 8.33 ml	wheat 1.0 mg	Phytosterol F 5.8 mg	23	25	28
Example 15	aqueous dispersion 3.13 ml	rice bran 1.0 mg	Phytosterol F 5.8 mg	25	28	30
Compara. Example 5	aqueous dispersion 1.45 ml	konjac 1.0 mg	-	20	24	26
Compara. Example 6	aqueous dispersion 1.45 ml	-	Phytosterol F 5.8 mg	10	13	16
Compara. Example 7	solution 1.45 ml	-	-	8	12	14

Phytosterol F: Plant sterols of soybean oil origin mainly comprising β -sitosterol, available from Eisai Co., Ltd.

Answers and scores:

4 ... The skin roughening was completely healed.

3 ... The skin condition was improved greatly.

2 ... The skin condition was improved.

5 1 ... The skin condition was unchanged.

0 ... The skin condition got worse.

Example 16: Preparation of cosmetic containing
sphingoglycolipids of konjac origin

10 An ethyl acetate:methanol (=9:1; 200 ml) mixed
solvent was passed through a column packed with 150 ml of
silica gel (Silica Gel 60, available from Nacalai tesque,
Inc.; 70 to 230 mesh). When the solvent liquid level
reached the upper edge of silica gel, and effusion became
15 slow, 1.9 g of precipitate A of Example 4 dissolved in
hexane to make 6.0 ml was introduced into the column. Then
400 ml of an ethyl acetate:methanol (=9:1) mixed solvent
was passed through, and the effluent was collected in 20 ml
fractions. Each fraction was analyzed by thin layer
20 chromatography. The 12th to 17th fractions developed deep
spots of sphingoglycolipids. The fractions in this range
were combined, and the solvent was removed by evaporation
to give 386 mg of a solid. HPLC revealed that the
sphingoglycolipids content of the solid was 226 mg,
25 indicating that the resulting sphingoglycolipids-containing
product had a sphingoglycolipids concentration of 58.5 wt%.

An ethyl acetate:methanol (=95:5) mixed solvent (200 ml) was passed through a column packed with 150 ml of silica gel (Silica Gel 60, available from Nacalai tesque, Inc.; 70 to 230 mesh). When the solvent liquid level reached the upper edge of silica gel, and effusion became slow, 386 mg of the solid prepared above by silica gel column chromatography and dissolved in a small amount of hexane was introduced into the column. Then 800 ml of an ethyl acetate:methanol (=95:5) mixed solvent was passed through, and the effluent was collected in 20 ml fractions. Each fraction was analyzed by thin layer chromatography. A single spot of sphingoglycolipids was observed with the 20th to 26th fractions. The fractions in this range were combined, and the solvent was removed by evaporation to give 87 mg of a solid. HPLC revealed that the sphingoglycolipids content of the solid was 86 mg, indicating that the resulting sphingoglycolipids-containing product had a sphingoglycolipids concentration of 98.9 wt%.

The solid (50.6 mg) was put into 10 ml of water with stirring and dispersed by ultrasonication using BRANSON 3200, available from Yamato Scientific Co., Ltd., for 30 minutes to obtain a cosmetic containing sphingoglycolipids of konjac tobiko powder origin according to the present invention. The resulting sphingoglycolipids-containing aqueous dispersion was uniform and produced no sediment when allowed to stand for

one day. This aqueous dispersion contained 50 mg of sphingoglycolipids of konjac tobiko powder origin.

Example 17: Preparation of cosmetic containing

5 sphingoglycolipids of potato origin

Five grams of the potato peel extract obtained in Example 3 was dissolved in 10.0 g of ethanol, and the solution was introduced into 100 g of water while stirring.

10 The mixture was further stirred while in a dispersed state. Thirty minutes later, 2.5 g of sodium chloride was added, and the stirring was continued for additional 10 minutes to give a light brown precipitate, which weighed 3.0 g. The sphingoglycolipids content was 0.18 g as measured by HPLC. After washing with water, the extract
15 had a sphingoglycolipids purity of 6.0 wt%. The washing operation was repeated once more to give 2.8 g of a precipitate, which was found to have a sphingoglycolipids content of 0.18 g as a result of HPLC. The purity of the sphingoglycolipids obtained after washing twice was

20 6.4 wt%. The resulting product containing sphingoglycolipids of potato peel origin was purified twice by silica gel column chromatography in the same manner as in Example 16 to obtain 38 mg of a sphingoglycolipids-containing product having a sphingoglycolipids content of
25 96.8 wt%.

1
2
3
4 The above-described series of operations were
5 repeated several times. The thus produced product
6 containing sphingoglycolipids of potato peel origin
7 (51.7 mg) was put into 10 ml of water while stirring and
8 dispersed by 30-minute ultrasonication by use of BRANSON
9 3200, available from Yamato Scientific Co., Ltd., to obtain
10 a cosmetic containing the sphingoglycolipids of potato peel
11 origin according to the present invention. The resulting
12 sphingoglycolipids-containing aqueous dispersion was
13 uniform and generated no sediment after one-day standing.
14 This aqueous dispersion contained 50 mg of
15 sphingoglycolipids of potato peel origin.

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
22

precipitate, which was found to have a sphingoglycolipids content of 0.04 g as a result of HPLC. The purity of the sphingoglycolipids obtained after washing twice was 1.2 wt%. The resulting product containing
5 sphingoglycolipids of wheat origin was purified twice by silica gel column chromatography in the same manner as in Example 16 to obtain 2 mg of a sphingoglycolipids-containing product having a sphingoglycolipids content of 97.5 wt%.

10 The above-described series of operations were repeated several times. The thus produced product containing sphingoglycolipids of wheat origin (51.3 mg) was put into 10 ml of water while stirring and dispersed by 30-minute ultrasonication by use of BRANSON 3200, available
15 from Yamato Scientific Co., Ltd. The resulting sphingoglycolipids-containing aqueous dispersion was uniform and generated no sediment after one-day standing. This aqueous dispersion contained 50 mg of sphingoglycolipids of wheat origin.

20 Comparative Example 9: Preparation of cosmetic containing sphingoglycolipids of rice bran origin

Ten grams of the defatted rice bran extract obtained in Comparative Example 2 was dissolved in 20.0 g
25 of ethanol, and the solution was introduced into 200 g of water while stirring. The mixture was further stirred

while in a dispersed state. Thirty minutes later, 5.0 g of sodium chloride was added, and the stirring was continued for 10 minutes to form a brown precipitate, which weighed 7.4 g. The sphingoglycolipids content of the precipitate was 0.24 g as measured by HPLC. After washing with water, the extract had a sphingoglycolipids purity of 3.2 wt%. The washing operation was repeated once more to give 6.9 g of a precipitate, which was found to have a sphingoglycolipids content of 0.22 g as a result of HPLC.

The purity of the sphingoglycolipids obtained after washing twice was 3.2 wt%. The resulting product containing sphingoglycolipids of rice bran origin was purified twice by silica gel column chromatography in the same manner as in Example 16 to obtain 28 mg of a sphingoglycolipids-containing product having a sphingoglycolipids content of 98.2 wt%.

The above-described series of operations were repeated several times. The thus produced product containing sphingoglycolipids of rice bran origin (50.9 mg) was put into 10 ml of water while stirring and dispersed by 30-minute ultrasonication by use of BRANSON 3200, available from Yamato Scientific Co., Ltd. The resulting sphingoglycolipids-containing aqueous dispersion was uniform and generated no sediment after one-day standing. This aqueous dispersion contained 50 mg of sphingoglycolipids of rice bran origin.

Test Example 5: Influences on SC water content

The skin SC water content improving effects of the cosmetics of the invention obtained in Examples 16 and 17 and the aqueous dispersions obtained in Comparative Examples 8 and 9 were tested with cooperation of ten female volunteers suffering from dry skin (five persons between the ages of 20 and 40 and five persons between the ages of 40 and 60). The test persons were asked to apply 0.5 ml of each aqueous dispersion on the inner side of the left upper arm for consecutive 30 days each. The skin SC water content of the inner side of the left upper arm was measured with Corneometer after 10 days, 20 days and 30 days. The results obtained are shown in Table 5 together with the results of control (the skin where the dispersion was not applied).

TABLE 5

	Initial Value	After 10 Days	After 20 Days	After 30 Days
Example 16 (tobiko powder origin)	47	65	75	78
Example 17 (potato origin)	48	63	71	73
Compara. Example 8 (wheat flour origin)	48	58	60	60
Compara. Example 9 (rice bran origin)	45	58	62	64
Control (no application)	49	45	47	46

It can be seen from Table 5 that applying sphingoglycolipids of konjac tobiko powder origin or potato peel origin produces excellent effects of improving SC water content of the skin.

Example 18 and Comparative Example 10: Preparation of sphingoglycolipids-containing beauty lotion

A beauty lotion containing sphingoglycolipids of konjac tobiko powder origin was prepared according to the formulation shown in Table 6 below (Example 18). A beauty lotion was prepared according to the same formulation except for the sphingoglycolipids (Comparative Example 10).

TABLE 6

Sphingoglycolipids-containing product of Example 16 (sphingoglycolipids concentration: 98.9%)	0.5 wt%
Propylene glycol	4.0 wt%
70% Sorbitol solution	3.0 wt%
Ethanol	5.0 wt%
Perfume	0.1 wt%
Methyl p-hydroxybenzoate	0.1 wt%
Sodium citrate	0.2 wt%
Purified water	balance

Test Example 6: Improvement on feel of use

The beauty lotions obtained in Example 18 and Comparative Example 10 were organoleptically tested by twenty females for evaluating the feel on application for 2 months. The results are shown in Table 7.

TABLE 7

	Duration of feel of moisture	Make-up Wear-ability	Make-up Dura-bility	Skin Elas-ticity	Skin Flexib-ility	Overall Judge-ment
Example 18	37	30	32	35	37	38
Compara. Example 10	28	28	31	26	26	27

Answers to questionnaire and scores:

2 ... Considerably good

1 ... Slightly good

0 ... Not good

The figures in Table 7 are total scores.

It is seen from Table 7 that the beauty lotion containing sphingoglycolipids of konjac tobiko powder origin exhibits excellent effects in duration of moisture
5 feel, skin elasticity and skin flexibility.

Industrial Applicability:

The sphingoglycolipids-containing functional foods and cosmetics according to the present invention contain
10 sphingoglycolipids, which exist in human skin and seem to play an important role in moisture retention, in a high concentration and produce excellent effects in improving the SC water content of human skin and improving rough
15 skin, atopic dermatitis, allergic dermatitis, pimples, and so forth. According to the production method according to the invention, sphingoglycolipids-containing functional foods and cosmetics can be produced easily and economically
20 by using as raw materials konjac tobiko powder, potato peel, oil cakes, etc. that have not been utilized as food at all.

While the invention has been described in detail and with reference to specific examples thereof, it will be apparent to one skilled in the art that various changes and
25 modifications can be made therein without departing from the spirit and scope thereof.

This application is based on Japanese Patent
Application Nos. 2000-219087 filed July 19, 2000, 2001-
021947 filed January 30, 2001, 2001-087695 filed March 26,
2001, and 2001-188393 filed June 21, 2001, which are

5 incorporated herein by reference.

2001-188393